

THE FIRST SUBSTELLAR SUBDWARF? DISCOVERY OF A METAL-POOR L DWARF WITH HALO KINEMATICS

ADAM J. BURGASSER^{1,2}, J. DAVY KIRKPATRICK³, ADAM BURROWS⁴, JAMES LIEBERT⁴, I. NEILL REID⁵, JOHN E. GIZIS⁶, MARK R. MCGOVERN¹, L. PRATO¹, & IAN S. MCLEAN¹

Accepted to ApJ

ABSTRACT

We present the discovery of the first L-type subdwarf, 2MASS J05325346+8246465. This object exhibits enhanced collision-induced H₂ absorption, resulting in blue NIR colors ($J - K_s = 0.26 \pm 0.16$). In addition, strong hydride bands in the red optical and NIR, weak TiO absorption, and an optical/J-band spectral morphology similar to the L7 DENIS 0205–1159AB imply a cool, metal-deficient atmosphere. We find that 2MASS 0532+8246 has both a high proper motion, $\mu = 2''.60 \pm 0''.15 \text{ yr}^{-1}$, and a substantial radial velocity, $v_{\text{rad}} = -195 \pm 11 \text{ km s}^{-1}$, and its probable proximity to the Sun ($d = 10\text{--}30 \text{ pc}$) is consistent with halo membership. Comparison to subsolar-metallicity evolutionary models strongly suggests that 2MASS 0532+8246 is substellar, with a mass of $0.077 \lesssim M \lesssim 0.085 M_{\odot}$ for ages 10–15 Gyr and metallicities $Z = 0.1 - 0.01 Z_{\odot}$. The discovery of this object clearly indicates that star formation occurred below the Hydrogen burning mass limit at early times, consistent with prior results indicating a flat or slightly rising mass function for the lowest-mass stellar subdwarfs. Furthermore, 2MASS 0532+8246 serves as a prototype for a new spectral class of subdwarfs, additional examples of which could be found in NIR proper motion surveys.

Subject headings: Galaxy: solar neighborhood — infrared: stars — stars: chemically peculiar — stars: individual (2MASS J05325346+8246465) — stars: low mass, brown dwarfs — subdwarfs

1. INTRODUCTION

Subdwarfs are metal-deficient stars, classically defined as lying below the stellar main-sequence in optical color-magnitude diagrams (Kuiper 1939). These objects are in fact not subluminescent but rather hotter (i.e., bluer in optical colors) than equivalent mass main-sequence dwarfs, a consequence of their reduced metal opacity (Chamberlin & Aller 1951; Sandage & Eggen 1959). Cool subdwarfs (spectral types sdK and sdM) are typically found to have halo kinematics ($\langle V \rangle = -202 \text{ km s}^{-1}$; Gizis 1997), and these objects are presumably relics of the early Galaxy, with ages of 10–15 Gyr. Because low mass subdwarfs have lifetimes far in excess of the age of the Galaxy, they are important tracers of Galactic chemical history, and are representatives of the first generations of star formation.

All of the coolest subdwarfs ($[\text{Fe}/\text{H}] \sim -1.2 \pm 0.3$) and extreme subdwarfs ($[\text{Fe}/\text{H}] \sim -2.0 \pm 0.5$; Gizis 1997) currently known have been identified in optical proper motion surveys, most notably Luyten's Half-Second Catalog (Luyten 1979a, LHS) and Two-Tenths Catalog (Luyten 1979b), the APM Proper Motion Survey (Scholtz et al. 2000), and the Galactic plane survey of Lepine, Shara, & Rich (2002). The substantial space velocities of halo subdwarfs allow them to stand out in these surveys amongst

the overwhelming multitude of similarly faint, but slowly moving, background stars. However, only a handful of very cool subdwarfs with spectral types sdM6/esdM6 and later⁷ have been found (Gizis 1997; Gizis & Reid 1997; Schweitzer et al. 1999; Lepine, Rich, & Shara 2003; Lepine, Shara, & Rich 2003). These stars exhibit characteristic spectral signatures of strong metal hydrides (CaH, MgH, AlH, FeH), weak and/or absent metal oxides (TiO, VO, CO), and enhanced collision-induced (CIA) H₂ absorption (Mould & Hyland 1976; Liebert & Probst 1987; Saumon et al. 1994; Leggett et al. 2000). With $T_{\text{eff}} \gtrsim 2900 \text{ K}$ (Leggett et al. 2000), the coolest subdwarfs known have masses just above the Hydrogen Burning Minimum Mass (HBMM)⁸, which ranges from $0.072 M_{\odot}$ for Solar composition ($Z = Z_{\odot}$) to $0.092 M_{\odot}$ for $Z = 0$ (Chabrier & Baraffe 1997; Burrows et al. 2001).

In contrast, hundreds of significantly cooler Population I, or disk, dwarfs have been identified⁹, predominately in the wide-field optical and near-infrared (NIR) surveys DENIS (Epchtein et al. 1997), 2MASS (Skrutskie et al. 1997), and SDSS (York et al. 2000). These discoveries include many dozens of substellar objects, and extend well beyond the M spectral class into the L (Kirkpatrick et al. 1999; Martín et al. 1999) and the T (Burgasser et al. 2002;

¹ Department of Physics & Astronomy, University of California at Los Angeles, Los Angeles, CA, 90095-1562; adam@astro.ucla.edu, mcgovern@astro.ucla.edu, lprato@astro.ucla.edu, mclean@astro.ucla.edu

² Hubble Fellow

³ Infrared Processing and Analysis Center, M/S 100-22, California Institute of Technology, Pasadena, CA 91125; davy@ipac.caltech.edu

⁴ Steward Observatory, University of Arizona, Tucson, AZ 85721; burrows@as.arizona.edu, liebert@as.arizona.edu

⁵ Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218; inr@stsci.edu

⁶ Department of Physics and Astronomy, University of Delaware, Newark, DE 19716; gizis@udel.edu

⁷ Based on the Gizis (1997) classification scheme.

⁸ The HBMM is defined here as the minimum mass for which core Hydrogen fusion balances luminosity at ages later than 10 Gyr.

⁹ A listing of known M, L, and T dwarfs is maintained by J. D. Kirkpatrick at <http://spider.ipac.caltech.edu/staff/davy/ARCHIVE/index.html>.

Geballe et al. 2002) classes. The success of recent surveys in identifying very cool dwarf star and brown dwarfs derives from their use of red-sensitive CCDs and infrared array detectors. As we proceed to cooler temperatures, the majority of emitted flux from both disk dwarfs and subdwarfs moves to NIR wavelengths. Unfortunately, proper motion surveys of sufficient temporal breadth to identify halo stars rely primarily on optical, typically photographic, imaging. Factoring in their intrinsic rarity (0.3% number density compared to disk dwarfs; Reid & Hawley 2000) and substantial evolution toward lower luminosities (Burrows et al. 1993; Chabrier & Baraffe 1997, see Figure 3), it is clear why few very cool subdwarfs have been found.

In this article, we present the discovery of a new cool subdwarf, identified in the 2MASS database. This object, 2MASS J05325346+8246465¹⁰, is a high motion source with a spectral morphology consistent with a metal deficient L dwarf. In § 2 we discuss the identification of 2MASS 0532+8246 and subsequent NIR and optical observations obtained at the Palomar 60" Telescope. In § 3 we describe spectroscopic observations of 2MASS 0532+8246 obtained with the Near-IR Spectrometer (McLean et al. 1998, 2000, hereafter NIRSPEC) and Low Resolution Imaging Spectrograph (Oke et al. 1995, hereafter LRIS) at Keck Observatory. In § 4 we derive the space motion of this object from the imaging and spectroscopic data. In § 5 we discuss the substellarity of 2MASS 0532+8246 based on subsolar metallicity models from Burrows et al. (2001). We discuss our results in § 6.

2. IDENTIFICATION OF 2MASS 0532+8246

2MASS 0532+8246 was identified in a search for T dwarfs using the 2MASS Working database (Burgasser et al. 2002). It was selected for its blue NIR colors ($J - K_s = 0.26 \pm 0.16$) and lack of an optical counterpart at the 2MASS position in both POSS-I and POSS-II photographic plates. A finder chart is given in Figure 1, and astrometric and photometric data from 2MASS are listed in Table 1. Follow-up NIR imaging using the Palomar 60" Infrared Camera (Murphy et al. 1995, hereafter IRCam) on 25 September 1999 (UT) determined that this object is not an uncatalogued minor planet. Gunn r - and z -band images obtained with the Palomar 60" Facility CCD Camera on 18 November 2002 and 25 January 2003 (UT) further confirmed the red optical/NIR colors of this object. Aperture photometry from 18 November 2002, calibrated with observations of the SDSS photometric standard BD +33° 4737 (Smith et al. 2002, correcting $(r', z') \rightarrow (\text{gunn } r, \text{gunn } z)$ using photometry from Kent 1985) yield $r > 20.9$ (5σ upper limit) and $z = 18.30 \pm 0.16$. As the NIR and optical/NIR colors of this object are consistent with those of other identified late-type L and T dwarfs (Kirkpatrick et al. 1999; Fan et al. 2000; Dahn et al. 2002), 2MASS 0532+8246 was initially believed to be a firm T dwarf candidate.

3. SPECTROSCOPIC OBSERVATIONS

3.1. NIR Spectrum

We obtained NIR spectra of 2MASS 0532+8246 on 24 December 2002 (UT) using NIRSPEC on the Keck-II 10m Telescope. Conditions were clear with seeing of 0".6. We observed the object in three spectral configurations, N3 ($1.14 - 1.38 \mu\text{m}$), N5 ($1.51 - 1.79 \mu\text{m}$), and N6b ($1.94 - 2.32 \mu\text{m}$), corresponding roughly to the J, H, and K NIR bands. Spectral resolution was $R \approx 2000$ for our 0".38 slit. For each observation, two exposures of 300s each were obtained, dithering 20" along the slit. All observations were made in the airmass range 2.6–3.0. Immediately after each sequence of exposures, internal flat field and NeAr arc lamps were observed for calibration, and the A0 star HD 34360 was observed just before or after the target source in each spectral order at similar airmasses. Data were reduced using the REDSPEC package¹¹; a detailed description of our reduction procedures is given in McLean et al. (2003). Final flux calibration of the individual spectral orders was done using 2MASS photometry; however, because the orders do not span the entire photometric bandpasses (Cutri et al. 2003), our final flux calibration may be somewhat underestimated at J-band, but reasonably correct for the H- and K-bands.

The resulting NIR spectrum is shown in Figure 2a. The $1.5 - 2.4 \mu\text{m}$ spectral energy distribution is fairly smooth and blue, reminiscent of an early-type main-sequence star or white dwarf, and quite unlike the much redder NIR spectra of late-type M and L dwarfs (cf. overlay of the L7 DENIS 0205–1159AB in Figure 2a; Delfosse et al. 1997; Kirkpatrick et al. 1999; McLean et al. 2003). However, the red optical/NIR colors and optical molecular bands (see below) of 2MASS 0532+8246 are inconsistent with a hot stellar atmosphere. We instead attribute the shape of the NIR spectrum to enhanced absorption by the CIA H_2 1-0 quadrupole band centered near $2.5 \mu\text{m}$ (Saumon et al. 1994; Borysow, Jørgensen, & Zheng 1997). This pressure-sensitive band, present in late-type L and T dwarfs (Toku-naga & Kobayashi 1999; Burgasser et al. 2002) and cool subdwarfs (Leggett et al. 2000), is generally broad and featureless, consistent with the observed spectrum. Strong H_2 absorption may explain the absence of CH_4 absorption bands at 1.6 and $2.2 \mu\text{m}$, the defining features of T dwarfs; and the $2.3 \mu\text{m}$ CO band, a key spectral feature in M, L, and early-type T dwarf spectra. On the other hand, H_2O absorption appears to be present at $1.3 - 1.5$ and $1.7 - 2.0 \mu\text{m}$, and minor features throughout the H-band are likely attributable to weaker H_2O lines. Two weak features at 1.573 and $1.626 \mu\text{m}$ are present, possibly attributable to FeH absorption (Cushing et al. 2003) or poor correction of telluric OH lines; these features are not coincident with the $1.6 \mu\text{m}$ CH_4 band. Overall, this portion of the spectrum of 2MASS 0532+8246 is quite unlike any known late-type M, L, or T dwarf, based on its blue slope and absence of CH_4 and CO bands.

The J-band spectrum, shown in detail in Figure 2b, is far more rich, with atomic lines of K I ($1.1690, 1.1773, 1.2543, 1.2522 \mu\text{m}$) and Fe I ($1.1528, 1.1886, 1.1976, 1.2128, 1.3210 \mu\text{m}$), and strong molecular bands of FeH ($1.194, 1.239$

¹⁰ Source designations for the 2MASS Point Source Catalog are given as "2MASS Jhhmmss[.jss±ddmmss[.s]". The suffix conforms to IAU nomenclature convention and is the sexagesimal R. A. and decl. at J2000 equinox. We adopt a shorthand notation of "2MASS hhmm±ddhh" throughout the remainder of this Letter.

¹¹ See <http://www2.keck.hawaii.edu/inst/nirspec/redspec/index.html>.

μm) and H_2O ($1.3 \mu\text{m}$) observed. Above the spectrum we plot an opacity spectrum of FeH (Dulick et al. 2003), which shows that much of the fine structure observed is caused by this molecule. There are no lines of Mg I, Ca I, or Al I present in this spectral region, and the ϕ TiO system around $1.25 \mu\text{m}$ (Jørgensen 1994) is either absent or obscured by FeH absorption. The overall spectral morphology at J-band is quite similar to the L7 DENIS 0205–1159AB, with the notable exception of stronger FeH and H_2O absorption.

3.2. Optical Spectrum

We obtained a red optical spectrum of 2MASS 0532+8246 on 3 January 2003 (UT) using LRIS on the Keck-I 10m Telescope. Conditions were clear with sub-arcsecond seeing. Two exposures of 1200 sec each were obtained through the red channel, dithered by $2''$ between exposures along the $1''$ (4.7 pixels) slit. We employed the 400 lines mm^{-1} grating blazed at $8500 \text{ \AA}$, yielding 6300–10100 $\text{\AA}$ spectra with $7 \text{ \AA}$ resolution ($R \sim 1200$). Dispersion on the chip was $1.9 \text{ \AA pixel}^{-1}$. The OG570 order-blocking filter was used to suppress higher-order light. Observations of the B1 V flux standard Hiltner 600 (Hamuy et al. 1994) were obtained for flux calibration, and the G0 V star HD 38847 was observed after the target at similar airmass (2.4) for telluric calibration. HeNeAr arc lamp exposures were taken immediately after the target observations for wavelength calibration, and quartz lamp flat-field exposures (reflected off of the interior dome) were observed at the start of the night to correct for detector response. Data reduction procedures, using standard IRAF¹² routines, are discussed in detail in Kirkpatrick et al. (2003).

The final flux-calibrated and telluric-corrected optical spectrum for 2MASS 0532+8246 is included in Figure 2a and shown in detail in Figure 2c. Strong metal hydride bands of FeH (8692, 9896 $\text{\AA}$), CrH (8611, 9969 $\text{\AA}$), and CaH (6750 $\text{\AA}$) are clearly discernable. The 9896 $\text{\AA}$ FeH Wing-Ford band is the strongest seen in any cool spectrum to date, and absorption from the shorter wavelength FeH and CrH bands extends to roughly 9100 $\text{\AA}$. Weaker bands of TiO (7053, 8432 $\text{\AA}$) and H_2O (9250, 9400 $\text{\AA}$) are also seen. We point out an absorption feature between 9570–9700 $\text{\AA}$ which is too strong to be attributable to intrinsic H_2O absorption, based on the weakness of the 9250 $\text{\AA}$ band; it is also not properly placed for FeH, CrH, TiH, or CH_4 absorption. This band is weakly present in late-type M and early-type L dwarf spectra, and may be a hitherto unrecognized metal hydride band. It is also possible, although unlikely, that this feature is attributable to residual telluric absorption, as telluric corrections were made only to 9650 $\text{\AA}$ (the extent of the strong terrestrial H_2O bands; Stevenson 1994). Individual alkali lines of Na I (8183, 8195 $\text{\AA}$), Rb I (7800, 7948 $\text{\AA}$), and Cs I (8521, 8943 $\text{\AA}$) are present, while the resonance K I doublet (7665, 7699 $\text{\AA}$) is exceptionally strong and pressure-broadened, as seen in late-type L and T dwarfs (Kirkpatrick et al. 1999; Liebert et al. 2000). The red optical/NIR colors of 2MASS 0532+8246 are clearly a consequence of the K I red wing; integrating the spectrum over the I_C bandpass (Bessell 1990) yields $I_C \approx 19.2$ and $I_C - J \approx 4.0$, similar to

mid- and late-type L dwarfs (Dahn et al. 2002). The blue end of the spectrum appears to be similarly suppressed by the pressure-broadened Na I doublet (5890, 5896 $\text{\AA}$). Both Li I (6708 $\text{\AA}$) absorption and $\text{H}\alpha$ (6563 $\text{\AA}$) emission lines are absent. The overall spectral morphology, excluding hydride bands and TiO, is again quite similar to the L7 DENIS 0205–1159AB, suggesting a rather cool atmosphere.

3.3. Characterization and Spectral Classification

2MASS 0532+8246 is not a T dwarf, based on the absence of the 1.6 and $2.2 \mu\text{m}$ CH_4 bands (Burgasser et al. 2002; Geballe et al. 2002), and the presence of strong FeH and CrH and weak TiO and Na I features in the optical (Burgasser, Kirkpatrick, & Liebert 2003). It appears that this object is instead a metal-deficient, late-type L dwarf. Enhanced CIA H_2 and strong hydride bands in the optical and NIR spectra of 2MASS 0532+8246 are consistent with a metal-poor atmosphere. Both are consequences of the overall reduced metal opacity, resulting in larger column abundances of the remaining chemical species and increased relative abundance of metal hydrides over double-metal species. The presence of the 7053 and 8432 $\text{\AA}$ TiO bands is somewhat at odds with this interpretation, particularly since TiO is generally weak or absent in disk dwarfs later than L5 (Kirkpatrick et al. 1999). One possibility is that the mechanism of Ti depletion in these objects – incorporation into multiple-metal condensates of CaTiO_3 , $\text{Ca}_3\text{Ti}_2\text{O}_7$, $\text{Ca}_4\text{Ti}_3\text{O}_{10}$, and Ti_2O_3 (Lodders 2002) – may be inhibited in the low-metallicity atmosphere of 2MASS 0532+8246. Chemical equilibrium calculations incorporating subsolar abundances are required to address this hypothesis. The TiO bands may also indicate a somewhat warmer atmosphere than the L7 comparison object DENIS 0205–1159AB.

Classifying 2MASS 0532+8246 is therefore not straightforward, particularly as there are no other metal-deficient L dwarfs currently known for direct comparison. The M subdwarf and extreme subdwarf sequences of Gizis (1997) terminate at sdM7 and esdM5.5, respectively, although recent late-type discoveries are classified via extrapolation and limited modifications of this scheme (Lepine, Rich, & Shara 2003; Lepine, Shara, & Rich 2003). Metal classification (d, sd, esd) in the Gizis scheme is based on the strength of the 7053 $\text{\AA}$ TiO band, which is exceedingly weak in the spectrum of 2MASS 0523+8246. Numerical classification is based on the 6750 $\text{\AA}$ CaH band, which is suppressed in this spectrum by Na I and K I absorption. Thus, a simple extrapolation of the Gizis scheme fails in the L dwarf regime. While there is general agreement between the spectral morphologies of 2MASS 0532+8246 and the DENIS 0205–1159AB, the optical/J-band hydride features and TiO bands are more similar to mid-type L dwarfs (Kirkpatrick et al. 1999; McLean et al. 2003). Additionally, there are no cool dwarf standards that match the $1.3\text{--}2.4 \mu\text{m}$ spectral region. We therefore tentatively classify this object a late-type sdL. Identification of additional metal-poor L subdwarfs may eventually allow the definition of a robust classification scheme.

¹² Image Reduction and Analysis Facility (IRAF) is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

4. SPACE MOTION

Examination of the IRCam and CCD followup images show that 2MASS 0532+8246 has moved significantly since it was first imaged by 2MASS. We therefore measured its proper motion by deriving its coordinates on each of the images, using the 2MASS coordinates of background stars for astrometric calibration. Results are listed in Table 2. A linear fit yields $\mu = 2''.60 \pm 0''.15 \text{ yr}^{-1}$ at position angle $130.0 \pm 1.8^\circ$, making 2MASS 0532+8246 one of the highest proper motion stars known (Bakos, Sahu, & Németh 2002). This object is not present in the LHS because of its optical faintness, although a faint counterpart may be present in the POSS-II R-band image at the extrapolated position (Figure 1). Residuals in the proper motion fit are indicative of parallactic motion, but positional uncertainties are too large for a reliable measure.

The radial velocity of 2MASS 0532+8246 was derived by cross-correlating its J-band spectrum with those of five early- and mid-type L dwarfs (chosen for their strong NIR FeH absorption) with measured radial velocities: 2MASS 0746+2000AB, 2MASS 1439+1929, G 196-3B, Kelu 1, and DENIS 1228-1547AB (Basri et al. 2000; Reid et al. 2002). All five of these objects have been observed as part of the NIRSPEC Brown Dwarf Spectroscopic Survey (McLean et al. 2003), and are therefore wavelength calibrated in the same manner as the data presented here. We derive a heliocentric $v_{rad} = -195 \pm 11 \text{ km s}^{-1}$, where the uncertainty is derived from the scatter of values amongst the comparison objects. This value is consistent with heliocentric velocity offsets in the $1.1690/1.1773 \mu\text{m}$ K I lines ($\langle v_{rad} \rangle = -175 \pm 17 \text{ km s}^{-1}$) and the optical Cs I and Rb I lines ($\langle v_{rad} \rangle = -192 \pm 24 \text{ km s}^{-1}$).

The substantial radial velocity and proper motion of 2MASS 0532+8246 identifies it as a high velocity source. Determining its total space motion, however, requires an estimate of its distance. If we assume 2MASS 0532+8246 is bound to the Galaxy (Galactocentric $v_{tot} \lesssim 500 \text{ km s}^{-1}$; Carney, Latham, & Laird 1988), we derive an upper limit distance of 54 pc assuming $(U, V, W)_\odot = (10.00, 5.25, 7.17) \text{ km s}^{-1}$ (Dehnen & Binney 1998) and Local Standard of Rest (LSR) velocity $V_{LSR} = 220 \text{ km s}^{-1}$ (Kerr & Lynden-Bell 1986). A spectrophotometric parallax estimate is highly uncertain, given the substantial redistribution of flux in the NIR and uncertainty in the numerical subtype for this object. However, assuming that the M_{IC} magnitude of 2MASS 0532+8246 is similar to that of mid- to late-type L dwarfs (Dahn et al. 2002, $17 \lesssim M_{IC} \lesssim 19$), we derive a conservative estimate of 10–30 pc. The corresponding heliocentric space velocities at the median distance are $(U, V, W) = (-13, -301, 28) \text{ km s}^{-1}$. The substantial V velocity, between -390 and -212 km s^{-1} over the adopted distance range, indicates zero or retrograde motion with respect to the LSR, and lends additional support that this object is a member of the halo population.

5. IS 2MASS 0532+8246 SUBSTELLAR?

Disk L dwarfs with spectral types later than $\sim \text{L4}$ are predominately substellar (Gizis et al. 2000); hence, the late-type spectral morphology of 2MASS 0532+8246 suggests that it too may be a brown dwarf. To test this hypothesis, we compared the estimated T_{eff} of this object to subsolar metallicity evolutionary models from Burrows

et al. (2001). In Figure 3, we plot T_{eff} versus mass for ages 10 and 15 Gyr and metallicities $Z = 0.1, 0.3$, and $0.01 Z_\odot$. At these late ages, the transition between stellar and substellar masses spans a substantial range in temperature, and is more dramatic for lower metallicities due to the reduced atmospheric opacities and hence enhanced luminosities (Burrows et al. 1993).

Deriving an accurate T_{eff} for 2MASS 0532+8246 is difficult without adequate distance or bolometric flux measurements. Solar-metallicity mid- to late-type L dwarfs have temperatures $1400 \lesssim T_{eff} \lesssim 1800 \text{ K}$ (Kirkpatrick et al. 1999; Basri et al. 2000; Pavlenko, Zapatero Osorio, & Rebolo 2000; Leggett et al. 2001; Stephens et al. 2001; Burgasser et al. 2002; Dahn et al. 2002). In general there is little difference between the temperatures of M-type subdwarfs and dwarf stars of the same numeric type (Gizis 1997; Leggett et al. 2000), although late-type M subdwarfs (e.g., the sdM7 LHS 377) may be 100–300 K hotter (Leggett et al. 2000). We therefore adopt a conservative temperature range of $1400 \lesssim T_{eff} \lesssim 2000 \text{ K}$. We further assume that the metallicity of 2MASS 0532+8246 lies in the range $0.01 \lesssim Z \lesssim 0.1 Z_\odot$, typical for sdM and esdM stars (Gizis 1997; Gizis & Reid 1997). For these parameters, we derive a mass range of $0.077 \lesssim M \lesssim 0.085 M_\odot$. Larger masses correspond to lower metallicities, so that in each case the derived mass is below the HBMM. Therefore, we conclude the 2MASS 0532+8246 is probably substellar, although more refined estimates of its atmospheric properties through further observations and spectral modelling are required to validate this result.

6. DISCUSSION

The existence of a substellar subdwarf is not unexpected, as mass functions for the nearby halo population (Gizis & Reid 1999) and globular clusters (Piotto & Zoccali 1999) are shallow but rising: $dN/dM \propto M^{-(0.5-1.3)}$ at $M \sim 0.15 M_\odot$, similar to the disk population (Reid et al. 1999; Chabrier 2002). However, the steep drop in T_{eff} , and hence luminosity, across the metal-poor substellar boundary implies that the vast majority of these halo brown dwarfs will have $T_{eff} \lesssim 1300 \text{ K}$; i.e., in the T dwarf regime and cooler (Burgasser et al. 2002). Indeed, one possible T-type subdwarf has already been identified, 2MASS 0937+2931 (Burgasser et al. 2002), an object which also exhibits unusually strong CIA H_2 and FeH absorption (Burgasser, Kirkpatrick, & Liebert 2003) and a substantial proper motion (Burgasser et al. 2003). Relatively few L-type subdwarfs should exist, however, as they span a much narrower range in mass at ages later than 10 Gyr. The discovery of 2MASS 0532+8246 may therefore be quite fortuitous.

Given its optical faintness, we expect that few counterparts to this late-type L subdwarf will be identified in current proper motion surveys. NIR proper motion surveys, on the other hand, could potentially reveal more of these objects by detecting them at the peak of their spectral energy distribution. A sizeable fraction of the sky ($\sim 30\%$) has been scanned more than once by 2MASS during its three years of observations, and analysis of these data may identify new, very cool high-motion stars. A second generation 2MASS survey could greatly improve on this work, adding substantially to the short list of very

cool subdwarfs currently known.

A. J. B. acknowledges useful discussions with K. Cruz and J. Wei-Chun during the preparation of the manuscript, and thanks our anonymous referee for her/his helpful criticisms. We are grateful for the assistance of our Keck Observing Assistants Joel Aycock and Gary Puniwai, and Instrument Specialist Paola Amico; and our Palomar Telescope Operators, Karl Dunscombe and Barrett “Skip” Staples. We also acknowledge the 2MASS staff for their laudable efforts on the 2MASS database, without which none of this research could have been possible. A. J. B. acknowledges support by NASA through Hubble Fellowship grant HST-HF-01137.01 awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., for NASA, under contract NAS 5-26555. A. S. B. acknowledges funding through NASA grants NAG5-10760 and NAG5-10629. Portions of the data presented herein were obtained at the W. M. Keck Observatory which is operated as a scientific

partnership among the California Institute of Technology, the University of California, and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W. M. Keck Foundation. This publication makes use of data from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center, funded by the National Aeronautics and Space Administration and the National Science Foundation. 2MASS data were obtained through the NASA/IPAC Infrared Science Archive, which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. POSS-I and POSS-II images were obtained from the DSS image server maintained by the Canadian Astronomy Data Centre, which is operated by the Herzberg Institute of Astrophysics, National Research Council of Canada. The authors wish to extend special thanks to those of Hawaiian ancestry on whose sacred mountain we are privileged to be guests.

REFERENCES

- Bakos, G. Á., Sahu, K. C., & Németh, P. 2002, *ApJS*, 141, 187
 Basri, G., Mohanty, S., Allard, F., Hauschildt, P. H., Delfosse, X., Martín, E. L., Forveille, T., & Goldman, B. 2000, *ApJ*, 538, 363
 Bessell, M. S. 1990, *PASP*, 102, 1181
 Borysow, A., Jørgensen, U. G., & Zheng, C. 1997, *A&A*, 324, 185
 Burgasser, A. J., Kirkpatrick, J. D., Liebert, J. 2003, *ApJ*, submitted
 Burgasser, A. J., Kirkpatrick, J. D., McElwain, M. W., Cutri, R. M., Burgasser, A. J., & Skrutskie, M. F. 2003, *AJ*, 125, 850
 Burgasser, A. J., et al. 2002, *ApJ*, 564, 421
 Burrows, A., Burgasser, A. J., Kirkpatrick, J. D., Liebert, J., Milsom, J. A., Sudarsky, D., & Hubeny, I. 2002, *ApJ*, 573, 394
 Burrows, A., Hubbard, W. B., Lunine, J. I., & Liebert, J. 2001, *Rev. of Modern Physics*, 73, 719
 Burrows, A., Hubbard, W. B., Saumon, D., & Lunine, J. I. 1993, *ApJ*, 406, 158
 Carney, B., Latham, D., & Laird, J. B. 1988, *AJ*, 96, 560
 Chabrier, G. 2002, *ApJ*, 567, 304
 Chabrier, G., & Baraffe, I. 1997, *A&A*, 327, 1039
 Chamberlin, J. W., & Aller, L. H. 1951, *ApJ*, 114, 52
 Cushing, M. C., Rayner, J. T., Davis, S. P., & Vacca, W. D. 2003, *ApJ*, 582, 1066
 Cutri, R. M., et al. 2003, http://www.ipac.caltech.edu/2mass/releases/allsky/doc/exp_hyp.html
 Dahn, C. C., et al. 2002, *AJ*, 124, 1170
 Dehnen, W., & Binney, J. J. 1998, *MNRAS*, 298, 387
 Delfosse, X., et al. 1997, *A&A*, 327, L25
 Dulick, M., Bauschlicher, C. W., Jr., Burrows, A., Sharp, C. M., Ram, R. S., & Bernath, P. 2003, *ApJ*, submitted
 Epchtein, N., et al. 1997, *The Messenger*, 87, 27
 Fan, X., et al. 2000, *AJ*, 119, 928
 Geballe, T. R., et al. 2002, *ApJ*, 564, 466
 Gizis, J. E. 1997, *AJ*, 113, 806
 Gizis, J. E., Monet, D. G., Reid, I. N., Kirkpatrick, J. D., Liebert, J., & Williams, R. 2000, *AJ*, 120, 1085
 Gizis, J. E., & Reid, I. N. 1997, *PASP*, 109, 849
 —. 1999, *AJ*, 117, 508
 Hamuy, M., Suntzeff, N. B., Heathcote, S. R., Walker, A. R., Gigoux, P., & Phillips, M. M. 1994, *PASP*, 106, 566
 Jørgensen, U. G. 1994, *A&A*, 284, 1791
 Kent, S. M. 1985, *PASP*, 97, 165
 Kerr, F. J., & Lynden-Bell, D. 1986, *MNRAS*, 221, 1023
 Kirkpatrick, J. D., et al. 1999, *ApJ*, 519, 802
 —. 2003, *AJ*, in preparation
 Kuiper, G. P. 1939, *ApJ*, 89, 549
 Leggett, S. K., Allard, F., Geballe, T., Hauschildt, P. H., & Schweitzer, A. 2001, *ApJ*, 548, 908
 Leggett, S. K., Allard, F., Dahn, C., Hauschildt, P. H., Kerr, T. H., & Rayner, J. 2000, *ApJ*, 535, 965
 Lepine, S., Rich, R. M., & Shara, M. M. 2003, *AJ*, in press (astro-ph/0209284)
 Lepine, S., Shara, M. M., & Rich, R. M. 2002, *AJ*, 124, 1190
 Lepine, S., Shara, M. M., & Rich, R. M. 2003, *ApJ*, 585, L69
 Liebert, J., & Probst, R. G. 1987, *ARA&A*, 25, 473
 Liebert, J., Reid, I. N., Burrows, A., Burgasser, A. J., Kirkpatrick, J. D., & Gizis, J. E. 2000, *ApJ*, 533, L155
 Lodders, K. 2002, *ApJ*, 577, 974
 Luyten, W. J. 1979a, *LHS Catalogue: A Catalogue of Stars with Proper Motions Exceeding 0'5 Annually* (Minneapolis: Univ. Minn. Press)
 —. 1979b, *New Luyten Catalogue of Stars with Proper Motions Larger than Two Tenths of an Arcsecond (NLTT)* (Minneapolis: Univ. Minn. Press)
 Martín, E. L., Delfosse, X., Basri, G., Goldman, B., Forveille, T., & Zapatero Osorio, M. R. 1999, *AJ*, 118, 2466
 McLean, I. S., Graham, J. R., Becklin, E. E., Figer, D. F., Larkin, J. E., Levenson, N. A., & Teplitz, H. I. 2000, *SPIE*, 4008, 1048
 McLean, I. S., McGovern, M. R., Burgasser, A. J., Kirkpatrick, J. D., Prato, L., & Kim, S. 2003, *ApJ*, submitted
 McLean, I. S., Prato, L., Kim, S. S., Wilcox, M. K., Kirkpatrick, J. D., & Burgasser, A. J. 2001, *ApJ*, 561, L115
 McLean, I. S., et al. 1998, *SPIE*, 3354, 566
 Møller, P., & Røed, A. R. 1976, *ApJ*, 208, 399
 Murphy, D. C., Persson, S. E., Pahre, M. A., Sivaramakrishnan, A., & Djorgovski, S. G. 1995, *PASP*, 107, 1234
 Oke, J. B., et al. 1995, *PASP*, 107, 375
 Pavlenko, Ya., Zapatero Osorio, M. R., & Rebolo, R. 2000, *A&A*, 355, 245
 Piotto, G., & Zoccali, M. 1999, *A&A*, 345, 485
 Reid, I. N., & Hawley, S. L. 2000, *New Light on Dark Stars* (Chichester: Praxis)
 Reid, I. N., Kirkpatrick, J. D., Liebert, J., Gizis, J. E., Dahn, C. C., & Monet, D. G. 2002, *AJ*, 124, 519
 Reid, I. N., et al. 1999, *ApJ*, 521, 613
 Sandage, A. R., & Eggen, O. J. 1958, *MNRAS*, 119, 278
 Saumon, D., Bergeron, P., Lunine, J. I., Hubbard, W. B., & Burrows, A. 1994, *ApJ*, 424, 333
 Scholtz, R.-D., Irwin, M., Ibata, R., Jahreiß, H., & Malkov, O. Yu. 2000, *A&A*, 353, 958
 Schweitzer, A., Shultz, R.-D., Stauffer, J., Irwin, M., & McCaughren, M. J. 1999, *A&A*, 350, L62
 Skrutskie, M. F., et al. 1997, in *The Impact of Large-Scale Near-IR Sky Surveys*, ed. F. Garzon (Dordrecht: Kluwer), p. 25
 Smith, J. A., et al. 2002, *AJ*, 123, 2121
 Stephens, D. C., Marley, M. S., Noll, K. S., & Chanover, N. 2001, *ApJ*, 556, L97
 Stevenson, C. C. 1994, *MNRAS*, 267, 904
 Tokunaga, A. T., & Kobayashi, N. 1999, *AJ*, 117, 1010
 York, D. G., et al. 2000, *AJ*, 120, 1579

TABLE 1
OBSERVATIONAL PROPERTIES OF THE LATE-TYPE SDL 2MASS 0532+8246.

$\alpha_{J2000}^{\text{a}}$	$5^h 32^m 53''.46$
$\delta_{J2000}^{\text{a}}$	$+82^\circ 46^m 46''.5$
I_C^{b}	19.2
Gunn z	18.30 ± 0.16
2MASS J	15.18 ± 0.06
2MASS H	14.90 ± 0.10
2MASS K_s	14.92 ± 0.15
μ	$2''.60 \pm 0''.15 \text{ yr}^{-1}$
θ	$130.0 \pm 1.8^\circ$
v_{rad}	$-195 \pm 11 \text{ km s}^{-1}$

^aEpoch 1 March 1999 (UT).

^bMeasured from optical spectrum; see § 3.2.

TABLE 2
ASTROMETRY OF 2MASS 0532+8246

Epoch (UT) (1)	Instrument (2)	α_{J2000} (3)	δ_{J2000} (4)	σ_α^{a} (5)	σ_δ^{a} (6)
01 Mar 1999 02:20	2MASS	$5^h 32^m 53''.46$	$+82^\circ 46^m 46''.53$	$0''.30$	$0''.30$
25 Sep 1999 11:53	P60 IRCAM	$5^h 32^m 54''.14$	$+82^\circ 46^m 45''.60$	$0''.07$	$0''.06$
18 Nov 2002 08:01	P60 CCD	$5^h 32^m 57''.37$	$+82^\circ 46^m 40''.33$	$0''.35$	$0''.23$
25 Jan 2003 06:33	P60 CCD	$5^h 32^m 57''.66$	$+82^\circ 46^m 40''.00$	$0''.37$	$0''.22$

^aUncertainty in (α, δ) derived from standard deviation of positions for background stars.




FIG. 1.— Finderchart for 2MASS 0532+8246, showing POSS-I (R-band), POSS-II (R-band), and 2MASS (J- and K_s-bands) fields. Images are scaled to the same spatial resolution, 5' on a side, with North up and East to the left. A 10'' box is centered at the position of the subdwarf in the 2MASS images and at the extrapolated position (assuming $\mu = 2''.6$ and $\theta = 130^\circ$) in the POSS-I and POSS-II images.




FIG. 2.— The spectrum of 2MASS 0532+8246 (thick black line) as compared to the L7 DENIS 0205–1159AB (thin grey line; data from Kirkpatrick et al. 1999; McLean et al. 2003). In all panels, spectral data for 2MASS 0532+8246 have been shifted by $v_{rad} = +195 \text{ km s}^{-1}$, and data for DENIS 0205–1159AB have been scaled to coincide at $1.27 \mu\text{m}$. Zero points are indicated by dotted lines. (a) Observed $0.63\text{--}2.35 \mu\text{m}$ spectrum, with NIRSPEC bands scaled to 2MASS photometry. Key atomic and molecular features are indicated; note that the $2.2 \mu\text{m}$ CH₄ and $2.3 \mu\text{m}$ CO bands present in the spectrum of DENIS 0205–1159AB (McLean et al. 2001) are not present in that of 2MASS 0532+8246. (b) *Top*: FeH absorption coefficient versus wavelength, from Dulick et al. (2003). *Bottom*: J-band spectrum of 2MASS 0532+8246, with line identifications for K I, Fe I, FeH, and H₂O. (c) Red optical spectrum, with key features indicated. Uncorrected telluric H₂O absorption in the DENIS 0205–1159AB data is indicated by the dashed bracket. Inset window shows a close-up of the $6350\text{--}7600 \text{ \AA}$ spectral region, highlighting strong CaH and weak TiO bands; no Li I or H α lines are seen.




FIG. 3.— T_{eff} versus mass for theoretical evolutionary models with $Z = 0.1$ (light grey), 0.03 (dark grey), and 0.01 ($Z_{\odot}$), and ages 10 (solid) and 15 (dashed) Gyr. HBMMs at 10 Gyr for all three metallicities are indicated by solid circles. Our adopted $1400 \lesssim T_{\text{eff}} \lesssim 2000$ K range for 2MASS 0532+8246 corresponds to masses of $0.077 \lesssim M \lesssim 0.085 M_{\odot}$ over all values of metallicity and age shown (hatched region). For each metallicity the derived mass is below the corresponding HBMM.